

A thermal theory for charged leptons

Ralf Hofmann

Institut für Theoretische Physik, Universität Heidelberg, Philosophenweg 16, 69120 Heidelberg, Germany

We propose that charged leptons are the ‘baryons’ of the gauge dynamics subject to a $SU(2)_e \times SU(2)_\mu \times SU(2)_\tau$ symmetry. The ‘mesons’ of this theory are neutral and thus candidates for dark matter. There is, in addition, a gauge group $SU(2)_{\text{CMB}}$ generating the CMB photon. We *compute* the value of α_{em} and the contribution to Λ_{cosmo} coming from $SU(2)_{\text{CMB}}$. We qualitatively explain the occurrence of dark matter, large intergalactic magnetic fields, a lepton asymmetry, the stability of T_{CMB} , and the deviation from thermal QED behavior of hot plasmas present in the interior of the sun and terrestrial nuclear fusion experiments.

PACS numbers: 12.38.Mh, 11.10.Wx, 12.38.G, 04.40.Nr

In [1] we have developed an analytic, inductive approach to hot $SU(N)$ pure gauge dynamics. Here we apply this approach. In the approach [1] we have observed that an $SU(2)$ gauge theory develops an electric phase (2nd order thermal phase transition), presumably at a temperature T_E^P tremendously higher than T_c (deconfinement temperature), maybe we have $T_E^P \sim M_P$ (M_P the Planck mass). In the electric phase $SU(2)$ is spontaneously broken to $U(1)$ by an adjoint Higgs field ϕ describing the condensation of calorons [1]. The tree-level massless (TLM) gauge boson - the photon - acquires mass by a one-loop radiative tadpole correction, see Fig. 1. This mass is given as

$$m_{\text{TLM}} = 0.433/(\sqrt{8}\pi)\sqrt{\Lambda_E^3/(2\pi T)} + O(1/e^2), \quad (1)$$

where Λ_E denotes the mass scale governing the electric phase, and e is the (large) $SU(2)$ gauge coupling. As a

function of T m_{TLM} has a global maximum at the critical temperature T_M^c where a mild 1st order thermal phase transition towards the magnetic phase takes place. The mass m_{TLM} decreases with $T^{-1/2}$ as T grows. Taking T to be room temperature, $T \sim 6.5 \times 10^{-2} \text{ eV}$, the temperature of visible radiation, $T \sim 6.5 \text{ eV}$, the temperature in the interior of the sun, $T \sim 10^7 \text{ eV}$ and assuming $\Lambda_E \sim 3.91 \times 10^{-4} \text{ eV}$ (see below) we arrive at a thermal photon mass in the electric phase of about $5.9 \times 10^{-7} \text{ eV}$, $5.9 \times 10^{-8} \text{ eV}$, $4.8 \times 10^{-11} \text{ eV}$, respectively. The *thermal* photon mass is not to be confused with the only valid bound on the photon mass ($\sim 10^{-14} \text{ eV}$) coming from precision measurements of the Coulomb potential [2, 3], see below. In unitary gauge off-diagonal gauge bosons ($V^\pm$) have large tree-level masses (TLH mode [1]) with small radiative corrections throughout the entire electric phase. There is *one* species of quasiclassical, stable BPS magnetic monopoles and antimonopoles in the electric phase [1, 4, 5] to which the massive gauge bosons ($V^\pm$) do not couple.

Under the intact $U(1)$ gauge symmetry in the electric phase the monopole charge g is given as $g = 2\pi/e$ [6, 7]. As the system cools down the electric coupling e strongly rises from a value very close to zero at T_E^P and then quickly relaxes to a value $e \sim 17.15$. Close to a temperature $T_M^c \ll T_E^P$ a regime of logarithmic blow-up is reached. At T_M^c $V^\pm$ gauge bosons decouple kinematically.

The value $e \sim 17.15$, see Fig. 2, is *independent* of the initial condition $e(T_E^P) = 0$ as long as T_E^P is sufficiently larger than T_M^c . Once the system has undergone the transition to the magnetic phase by condensing monopoles, the left-over photon is *exactly* massless [1] on the magnetic side of the phase boundary. In the magnetic phase the gauge coupling g and the photon mass quickly increases with decreasing temperature, see Fig. 1. The ground state of the system becomes *superconducting* with respect to the spontaneously broken $U(1)$ gauge symmetry. Considerably far away from the transition center vortices form due to a growing size of monopoles in the condensate [1]. In the magnetic phase, single crossings of center vortices (topological charge 1/2) have charge $2\pi/g$ with respect to the massive photon [10].




FIG. 1: Radiative corrections arising in the electric phase if gauge-boson fluctuations are assumed to be noninteracting thermal quasiparticles. (a) - (c) thermal mass corrections and (d) - (g) corrections to the vacuum pressure. Thin lines are associated with tree-level massless (TLM) modes and thick ones with tree-level heavy (TLH) modes.




FIG. 2: The gauge couplings in the electric and in the magnetic phase as a function of dimensionless temperature $\lambda_{E,M}$, see [1]. The ‘nervous’ behavior in the electric case is due to numerical interpolation errors. The plateau value of e can be read off at the downward peaks. It is $e = 17.15$.

At the critical temperature $T_M^c < T_E^c$ the magnetic gauge coupling g diverges logarithmically: photons decouple kinematically, and the system condenses the center vortices. A 1st order nonthermal phase transition takes place [1]. There are stable ‘hadrons’ with what we call one unit of electric charge in the Standard Model: ‘baryons’. Moreover, there is a stable neutral ‘hadron’, see Fig. 3. Considering *naively* only a single $SU(2)$ gauge group, the value of the electric charge is $g = 2\pi/e \sim 0.3664$. This would yield $\alpha_{em} = g^2/(4\pi) \sim 1/93.6$: a value which is considerably off the measured value $\alpha_{em} \sim 1/137$. So what went wrong in our naive approach to the electric charge of a lepton?

We propose that the CMB photon and the spectrum of charged leptons is described by gauge dynamics subject to the symmetry $SU(2)_{CMB} \times SU(2)_e \times SU(2)_\mu \times SU(2)_\tau$ (CMB stands for cosmic microwave background). The choice of $SU(2)$ factors derives from the experimental fact that no mass-degenerate charged leptons exist in nature, see below. The scale of the first factor is set by the CMB temperature $T_{CMB} = 2.728 \text{ K} \sim 2.182 \times 10^{-4} \text{ eV}$, the scales of the other factors are set by the respective masses of the charged leptons. The gauge dynamics subject to the $SU(2)_e \times SU(2)_\mu \times SU(2)_\tau$ is in the confining phase: no asymptotic gauge bosons exist here and the surviving, effective symmetry is a spontaneously broken *local* $Z_2^e \times Z_2^\mu \times Z_2^\tau$ symmetry [1]. Charged leptons are the ‘baryons’ of these gauge theories, ‘mesons’ are neutral and much less massive, see Fig. 3. The ‘mesons’ are candidates for dark matter, an interpretation of these particles is given in [8]. In [9] we have shown that a local Z_2 symmetry favors Fermi-Dirac statistic over Bose-Einstein statistics by a factor two if we assume the thermalization of the respective stable ‘hadronic’ states. In particular, the ‘baryons’ electron, muon, and tau are rendered fermions as a result of genuinely local Z_2 transformations.

The dynamics subject to $SU(2)_{CMB}$ is nonconfining. This theory is in its magnetic phase close to the electric phase boundary, where the photon is almost massless, see below and Fig. 4. At the temperature T_{CMB} the CMB photon is *quantum entangled* with

the (infinitely) heavy photons of the gauge theory with $SU(2)_e \times SU(2)_\mu \times SU(2)_\tau$ symmetry due to their mixing during an early epoch in the evolution of the Universe where the temperature was higher than the muon but lower than the pion mass.

Using this input, we now *derive* the value of α_{em} . Since the τ lepton has a very small life-time ($\delta t_\tau / \delta t_\mu \sim 10^{-7}$ but $m_\mu / m_\tau = 6 \times 10^{-2}$, decay by electroweak interactions) the photons of $SU(2)_\tau$ scattered by this particle were most probably never thermalized. Moreover, the gauge dynamics due to $SU(2)_\tau$ is contaminated by strong interactions for temperatures larger than the pion mass. But we only want to consider pure electromagnetism here. At $m_\mu < T < m_\pi$ we thus consider the gauge group $SU(2)_{CMB} \times SU(2)_e \times SU(2)_\mu$ only. We then have three species of thermalized photons a_μ^i ($i = \text{CMB}, e, \mu$) in the plasma which mix quantum mechanically by a thermal mass matrix (recall, that the tree-level massless modes acquire a mass by radiative corrections in the electric phase, Fig. 1 and (1)). In a thermal state it is very reasonable to assume that photon wave functions $\tilde{a}_\mu^i$ are generated by maximal mixing of the photon states originating from the respective dynamics of each $SU(2)$ factor, for example

$$\tilde{a}_\mu^{CMB} = 1/\sqrt{3}(a_\mu^{CMB} + a_\mu^e + a_\mu^\mu). \quad (2)$$

and similar for the orthogonal superpositions $\tilde{a}_\mu^e$ and $\tilde{a}_\mu^\mu$. When the temperature falls below m_μ , an epoch of exponential cosmological inflation of scale m_μ , which is terminated by a nonthermal 1st order transition, renders the $SU(2)_\mu$ gauge theory confining. The photon a_μ^μ decouples kinematically and is almost instantaneously removed from (2) as a fluctuating degree of freedom. This changes the wave function normalization of the remaining two photons by a factor $\sqrt{2}/\sqrt{3}$. The new maximal-mixing superposition still sees the charge of the muon because of a quantum entanglement with the kinematically decoupled photon a_μ^μ . The coherent superposition of a_μ^{CMB} and a_μ^e is now normalized as

$$\tilde{a}_\mu^{CMB} = 1/\sqrt{2} \left[\sqrt{2}/\sqrt{3}(a_\mu^{CMB} + a_\mu^e) \right]. \quad (3)$$

The further removal of a_μ^μ from the thermal spectrum by a period of exponential inflation of scale $T \sim m_e$ does not imply a new normalization since only the CMB photon survives kinematically. In the electric phase and away from the boundary to the magnetic phase monopoles are much heavier than the photons, $m_{\text{mon}}/m_{\text{photon}} = 64\pi^2/(\sqrt{2}e) = 25.5$. Shortly before the transition to the magnetic phase they experience a strong drop in their mass and condense subsequently in the magnetic phase. This condensation slowly rotates a superposition of monopoles belonging to different $SU(2)$ factors to the monopole of the $SU(2)$ factor which undergoes the electric-magnetic transition (monopole excitations, which disappear in the magnetic phase, re-appear as charged leptons in the center phase). We conclude,

that the above argument for photons does not apply to monopoles. The factor $\sqrt{2}/\sqrt{3}$ in (2) reduces the wave function of the original photon or, equivalently, the gauge coupling $g \rightarrow g_r = \sqrt{2}/\sqrt{3}g$. As a consequence, the real value of α_{em} is

$$\alpha_{em} = g_r^2/(4\pi) \sim 1/140.433. \quad (4)$$

The deviation from the measured value $\alpha_{em} \sim 1/137$ is 2.4%! We attribute this small mismatch to the omission of radiative corrections in our evolution of the gauge coupling e in the electric phase (recall that there is a *zeroth-order* in $1/e^2$ radiative correction to the photon mass). The result (4) is an impressive experimental confirmation for the validity of the analytic approach to thermal SU(N) Yang-Mills theory in [1]. The reader may wonder why we could predict the value of α_{em} from an analysis in the *electric* phase while charged leptons, ‘baryons’ of $SU(2)_e \times SU(2)_\mu \times SU(2)_\tau$, are solitons in the *center* phase of this theory. The solution to this apparent puzzle is indicated in Fig. 3 and explained in its caption. The electric charge is quantized by the single monopole residing in a ‘baryon’ (or charged lepton) of spin 1/2 as the crossing point of two center vortices. This ‘baryon’ has precisely the same charge as a monopole in the electric phase due to the topological equivalence of a center-vortex crossing and an isolated monopole [14].

In the magnetic phase monopoles condense, the magnetic coupling g is precisely zero at the magnetic-electric transition and the electric coupling e is infinite. The critical temperature T_E^c is the only point where a magnetic description is precisely dual to an electric description of the underlying SU(2) gauge theory. In the course of the magnetic-center transition, when center vortices condense, monopoles re-appear on the crossings of center vortices. In a given spatial region, charged lepton number is not conserved. This leads us to a new mechanism for the generation of lepton asymmetry. Recall Sakharov’s conditions [11] for this: (i) lepton number violation, (ii) violation of thermal equilibrium, and (iii) a CP violation in the interaction. We just discussed the violation of electric charge (in the sense of the Standard Model) during the magnetic-center transition. So condition (i) is met. The transition to the center phase is truly 1st order. It is preceded by an epoch of exponential inflation (a photon decouples kinematically shortly before the transition and thus the energy density ρ_{vac} is the one of the ground state, the equation of state is $P_{vac} = -\rho_{vac}$). So condition (ii) is met. We know that weak interactions violate CP due to a 3×3 CKM matrix for the three visible charged leptons. So there is a CP violating phase, condition (iii) is also met. The magnitude of lepton asymmetry depends on the mass of the charged lepton since this sets the scale for the strength of the 1st order transition to the center phase. It also depends on the structure of the CKM matrix. Lepton asymmetry should then be computable from the measured CKM matrix in the Standard Model and the possibility to estimate the strength of the 1st order transition. Let us now de-




FIG. 3: The evolution of an electric charge (in the sense of the Standard Model) through the three phases of an SU(2) gauge theory. The dots indicate monopoles. The thin (double) lines in the magnetic (center) phase denote (fat) center vortices. The ‘meson’ in the center phase is not charged under $U(1)_{em}$. It is a dark-matter candidate. In the presence of the asymptotic CMB photon there are no other stable ‘baryonic’ states (more than one monopole or antimonopole in a ‘baryon’ would either repel or annihilate).

rive a boundary condition for the gauge theory $SU(2)_{CMB}$ from experimental information. On cosmological scales we do not see neither center vortices nor do we see light-mass electric charges at any temperature in laboratory experiments [12] (monopoles and antimonopoles are extremely dilute in the electric phase). We do, however, see a massless photon in the CMB radiation which has an (almost) perfect thermal spectrum. This leads us to the conclusion that the CMB temperature T_{CMB} is slightly lower than the critical temperature T_E^c of the electric-magnetic transition. Only for $T_{CMB} + 0 = T_E^c$ do we have a *condensate* of monopoles (no explicit monopoles!) and not yet an explicit center vortex in our theory. Moreover, in our theory the CMB photon is exactly massless at $T_{CMB} = T_E^c - 0$ due to a logarithmic blow-up of e on the electric side of the phase boundary rendering $g = 0$. As a consequence, the Abelian Higgs mechanism is not operative close to the phase boundary. So the boundary condition is: $T_{CMB} = T_E^c$.

The cooling of the ground state into superconductivity for $T < T_{CMB}$ will change our universe drastically. The onset of this regime is already seen in the existence of *large intergalactic magnetic fields*. The generation of these magnetic fields is driven by one-sided temperature fluctuations which drive the associated, intergalactic region away from the electric-magnetic phase boundary into the magnetic phase, see Fig. 4. What is the mechanism that prevents us from running deep into the superconducting regime? The thermal mass of the CMB photon in the electric phase jumps from $m_{TLM} = 0.433/(\sqrt{8}\pi)\sqrt{\Lambda_E^3/(2\pi T_{CMB})} \sim 10^{-5}$ eV to zero across the phase boundary. In the magnetic phase the CMB photon acquires mass as temperature drops. As a function of temperature this creates a potential for the photon mass with a wall of height $\sim 10^{-5}$ eV to the right and smoothly increasing mass to the left of the minimum at $T = T_{CMB}$, see Fig. 4. This effect keeps the CMB




FIG. 4: Photon-mass dependence on temperature.

temperature stable for about another present age of the universe. The energy needed for gravitational expansion is taken from the ground state. The stabilizing effect is actually seen in a lattice simulation of the energy density ρ as a small dip of ρ shortly before the deconfinement transition [13]. In the far-distant future we will have to cope with increasing magnetic fields and magnetic vortices in our universe until a phase of exponential inflation will put to an end the existence of the CMB photon.

Since the masses of charged leptons are much higher than T_{CMB} ($m_e = 5 \times 10^5 \text{ eV} \sim 2.3 \times 10^9 T_{\text{CMB}} \sim \Lambda_e$) the dynamics of $SU(2)_e$, $SU(2)_\mu$, and $SU(2)_\tau$ are all in their confining phase at T_{CMB} . The vacuum energy densities and the vacuum pressures in all three theories are *precisely* zero [1] (this is protected by local Z_2 symmetries). Let us now calculate the electromagnetic, homogeneous contribution to the cosmological constant. At the electric-magnetic phase boundary we have $\lambda_M(0) = (1/4)^{1/3} \times 9.193$ where due to our above boundary condition $\lambda_M(0) \equiv 2\pi T_{\text{CMB}}/\Lambda_M$ and Λ_M denotes the magnetic scale of $SU(2)_{\text{CMB}}$, see [1]. Prescribing $T = T_{\text{CMB}} = 2.728 \text{ K} = 2.1824 \times 10^{-4} \text{ eV}$, yields $\Lambda_M = 2.368 \times 10^{-4} \text{ eV}$. The resulting vacuum energy density is $1/2 V_M = \pi T_{\text{CMB}} \Lambda_M^3 = (3.1 \times 10^{-4} \text{ eV})^4$. This is smaller than the commonly accepted value $\Lambda_{\text{cosmo}} \sim (10^{-3} \text{ eV})^4$. There are two possible reasons for this. First, we have used a tree-level analysis to determine the critical value T_E^c in our approach. Including radiative corrections in

the running of the electric gauge coupling e should yield a slightly smaller value than $\lambda_E^c = 9.193$ since massive TLM modes slow down the evolution, see [1]. This should imply a slightly higher value of $1/2 V_M$. Second, there are local regions with negative pressure induced by high-energy particle collisions subject to $SU(N)$ gauge dynamics. After coarse-graining this effect also contributes to Λ_{cosmo} .

Let us now discuss the validity of thermal QED. The additional photon belonging to $SU(2)_e$ can not be seen in terrestrial experiments with hot plasmas up to $T \sim m_e = 5 \times 10^5 \text{ eV}$. The highest temperatures reached in nuclear fusion experiments are about $T \sim 10^7 \text{ eV}$ - the temperature in the interior of the sun. Due to the occurrence of a second photon, the strong dilution of the electron density in the electric phase of $SU(2)_e$, and a considerable contribution of vacuum energy these plasmas should show a strong deviation from standard thermal QED behavior. All solar models rely on standard thermal QED and thus should be revised. This should have a major impact on the prediction of neutrino fluxes. As for Tokamaks we predict a strong increase of the magnetic field at temperatures close to the electron mass and a disappearance of electronic electric charge at about 1.8 times this temperature. This effect could be measured by applying a large and homogeneous electric field to the Tokamak. Positively charged ions should then move to one end and no negative charge should be measured at the other end of the Tokamak.

To summarize, we have proposed a strongly interacting gauge theory underlying electrodynamics, and we have explored some of the consequences of this theory. The apparent structurelessness of the electron and the muon, as it is seen in high-energy experiments for the modulus of momentum transfers away from the electron and muon mass, is explained in [8].

Acknowledgments: It is a pleasure to thank Qaisar Shafi and Zurab Tavartkiladze for very useful discussions and Pierre van Baal for a correspondence. The hospitality and financial support of CERN's theory division are thankfully acknowledged.

-
- [1] R. Hofmann, hep-ph/0312046.
 - [2] E. Adelberger, G. Dvali, and A. Gruzinov, hep-ph/0306245.
 - [3] E. R. Williams, J. E. Faller, and H. A. Hill, Phys. Rev. Lett. **26**, 721 (1971).
 - [4] W. Nahm, Lect. Notes in Physics. 201, eds. G. Denaro, e.a. (1984) p. 189.
 - [5] T. C. Kraan and P. van Baal, Nucl. Phys. B **533**, 627 (1998); T. C. Kraan and P. van Baal, Phys. Lett. B **435**, 389 (1998).
 - [6] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
 - [7] P. Hasenfratz and G. 't Hooft, Phys. Rev. Lett. **36**, 1119 (1976).
 - [8] R. Hofmann, hep-ph/0401017.
 - [9] R. Hofmann, hep-ph/0312051.
 - [10] H. Reinhardt, O. Schroder, T. Tok, and V. Ch. Zhukovsky, Phys. Rev. D **66**, 085004 (2002).
 - [11] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5**, 32 (1967); JETP Lett. **5**, 24 (1967); Sov. Phys. Usp. **34**, 392 (1991); Usp. Fiz. Nauk **161**, 61 (1991).
 - [12] This charged particle can not be seen in terrestrial experiments because, locally, thermal equilibrium is always violated on scales much larger than T_{CMB} . The temperature of liquid ^3He may be lowered to milli-Kelvins but this temperature is the temperature of heavy fermionic compounds with eV electrons residing in their interiors. I thank Paul Clegg for a discussion of this point.
 - [13] J. Engels *et al.*, Nucl. Phys. B **205**[FS5], 545 (1982).

[14] The continuous variation of the magnetic charge in dependence of temperature in the magnetic phase is due to the occurrence of a monopole-antimonopole *condensate*,

that is, the percolation of single (anti)monopole world-lines.